

Package leaflet: Information for the user

Omeflex special without electrolytes emulsion for infusion

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Omeflex special without electrolytes is and what it is used for
2. What you need to know before you use Omeflex special without electrolytes
3. How to use Omeflex special without electrolytes
4. Possible side effects
5. How to store Omeflex special without electrolytes
6. Contents of the pack and other information

1. What Omeflex special without electrolytes is and what it is used for

Omeflex special without electrolytes contains fluids and substances called amino acids and fatty acids that are essential for the body to grow or to recover. It also contains calories in the form of carbohydrates and fats.

Omeflex special without electrolytes is given to adults, adolescents and children older than two years.

You are given Omeflex special without electrolytes when you are unable to eat food normally. There are many situations when this might be the case, for example when you are recovering from surgery, injuries, or burns, or when you are unable to absorb food from your stomach and gut.

2. What you need to know before you use Omeflex special without electrolytes

Do not use Omeflex special without electrolytes

- if you are allergic to any of the active substances, to egg, peanut, soybean or fish or to any of the other ingredients of this medicine (listed in section 6).
- This medicine must not be given to newborn infants, infants and toddlers under two years old.

Also, do not use Omeflex special without electrolytes if you suffer from any of the following:

- life-threatening blood circulation problems such as those that can occur if you are in a state of collapse or shock
- heart attack or stroke
- severely impaired blood clotting function, bleeding risk (severe coagulopathy, aggravating haemorrhagic diatheses)

- blocking of blood vessels by blood clots or fat (embolism)
- severe liver failure
- impaired bile flow (intrahepatic cholestasis)
- severe kidney failure in the absence of kidney replacement therapy
- disturbances of your body salt composition
- fluid deficit or excess water in your body
- water on your lungs (pulmonary oedema)
- severe heart failure
- certain metabolic disorders such as
 - too much lipid (fat) in the blood
 - inborn errors of amino acid metabolism
 - abnormally high blood sugar level that needs more than 6 units of insulin per hour to be controlled
 - abnormalities of metabolism that may occur after operations or injuries
 - coma of unknown origin
 - insufficient supply of oxygen to tissues
 - abnormally high acid level in the blood.

Warnings and precautions

Talk to your doctor, pharmacist or nurse before using Omeflex special without electrolytes.

Please inform your doctor if:

- you have heart, liver or kidney problems
- you suffer from certain types of metabolic disorders such as diabetes, abnormal blood fat values and disorders of your body fluid and salt composition or your acid-base balance

You will be monitored closely to detect early signs of an allergic reaction (such as fever, shivering, rash, or shortness of breath) when you receive this medicine.

Further monitoring and tests such as various examinations of blood samples will be applied to make sure that your body handles the administered foodstuffs properly.

This medicine is an electrolyte free solution. Your healthcare professional will also take measures to ensure that your body's fluid and electrolyte requirements are met. In addition to Omeflex special without electrolytes you may receive further nutrients (foodstuffs) in order to fully cover your requirements.

Children

This medicine must not be given to newborn infants, infants and toddlers under two years old.

Other medicines and Omeflex special without electrolytes

Tell your doctor, pharmacist or nurse if you are taking, have recently taken or might take any other medicines.

Omeflex special without electrolytes can interact with some other medicines. Please tell your doctor, pharmacist or nurse if you are taking or receiving any of the following:

- insulin
- heparin
- medicines that prevent undesirable blood clotting such as warfarin or other coumarin derivatives

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. If you are pregnant you will receive this medicine only if the doctor or pharmacist considers it absolutely necessary for your recovery. There is no data available about the use of Omeflex special without electrolytes in pregnant women.

Breast-feeding is not recommended for mothers on parenteral nutrition.

Driving and using machines

This medicine is normally given to immobile patients, e.g. in a hospital or clinic which would exclude driving or using machines. However, the medicine itself has no effect on the ability to drive and use machines.

Omeflex special without electrolytes contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per multichamber bag, i.e. it is essentially 'sodium-free'.

3. How to use Omeflex special without electrolytes

This medicine is administered by intravenous infusion (drip), that is, through a small tube directly into a vein. This medicine will be administered through one of your large (central) veins only. The recommended duration of infusion for a parenteral nutrition bag is maximum 24 h.

Your doctor or pharmacist will decide how much of this medicine you need and for how long you will require treatment with this medicine.

Use in children

This medicine must not be given to newborn infants, infants and toddlers under 2 years old.

Your doctor will decide how much of this medicine your child needs and for how long your child will require treatment with this medicine.

If you use more Omeflex special without electrolytes than you should

If you have received too much of this medicine, you may suffer from a so-called 'overload syndrome' and the following symptoms:

- fluid excess and electrolyte disorders
- water on your lungs (pulmonary oedema)
- loss of amino acids through the urine and disturbed amino acid balance
- vomiting, feeling sick
- shivering
- high blood sugar level
- glucose in the urine
- fluid deficit
- blood much more concentrated than normal (hyperosmolality)
- impairment or loss of consciousness due to extremely high blood sugar
- enlargement of the liver (hepatomegaly) with or without jaundice (icterus)
- enlargement of the spleen (splenomegaly)

- fat deposition in the inner organs
- abnormal values of liver function tests
- reduction of red blood cell count (anaemia)
- reduction of white blood cell count (leucopenia)
- reduction of blood platelet count (thrombocytopenia)
- increase of immature red blood cells (reticulocytosis)
- rupture of blood cells (haemolysis)
- bleeding or a tendency to bleeding
- impairment of blood coagulation (as can be seen by changes of bleeding time, coagulation time, prothrombin time etc.)
- fever
- high blood fat levels
- loss of consciousness

If any of these symptoms occur, the infusion must be stopped immediately.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The following side effects may be serious. If any of the following side effects occur, tell your doctor immediately, he will stop giving you this medicine:

Rare (may affect up to 1 in 1,000 people):

- allergic reactions, for example skin reactions, shortness of breath, swelling of the lips, mouth, and throat, difficulty breathing

Other side effects include:

Uncommon (may affect up to 1 in 100 people):

- feeling sick, vomiting, loss of appetite

Rare (may affect up to 1 in 1,000 people):

- increased tendency for your blood to clot
- bluish discolouration of the skin
- shortness of breath
- headache
- flushing
- reddening of skin (erythema)
- sweating
- chills
- feeling cold
- high body temperature
- drowsiness
- pain in the chest, back, bones or lumbar region
- decrease or increase in blood pressure

Very rare (may affect up to 1 in 10,000 people):

- abnormally high blood fat or blood sugar values
- high levels of acidic substances in your blood
- Too much lipid can lead to fat overload syndrome, for more information on this please see under the heading “If you use more Omeflex special without electrolytes than you should” in section 3. Symptoms normally disappear when the infusion is stopped.

Not known (frequency cannot be estimated from the available data):

- reduction of white blood cell count (leucopenia)
- reduction of blood platelet count (thrombocytopenia)
- impaired bile flow (cholestasis)

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Omeflex special without electrolytes

Keep this medicine out of the sight and reach of children.

Do not store above 25 °C.

Do not freeze. If accidentally frozen, discard the bag.

Keep the bag in the protective overwrap in order to protect from light.

Do not use this medicine after the expiry date which is stated on the label. The expiry date refers to the last date of that month.

6. Contents of the pack and other information

What Omeflex special without electrolytes contains

The active substances in the ready-for-use mixture are:

<i>from the top chamber (glucose solution)</i>	in 1000 ml	in 625 ml	in 1250 ml	in 1875 ml
Glucose monohydrate	158.4 g	99.00 g	198.0 g	297.0 g
equivalent to glucose	144.0 g	90.00 g	180.0 g	270.0 g

<i>from the middle chamber (fat emulsion)</i>	in 1000 ml	in 625 ml	in 1250 ml	in 1875 ml
Medium-chain triglycerides	20.00	12.5	25.0	37.5
Soya-bean oil, refined	16.00	10.0	20.0	30.0
Omega-3-acid triglycerides	4.00	2.500	5.00	7.50

<i>from the bottom chamber (amino acid solution)</i>	in 1000 ml	in 625 ml	in 1250 ml	in 1875 ml
Isoleucine	3.284 g	2.053 g	4.105 g	6.158 g

Leucine	4.384 g	2.740 g	5.480 g	8.220 g
Lysine monohydrate equivalent to lysine	3.576 g 3.184 g	2.235 g 1.990 g	4.470 g 3.979 g	6.705 g 5.969 g
Methionine	2.736 g	1.710 g	3.420 g	5.130 g
Phenylalanine	4.916 g	3.073 g	6.145 g	9.218 g
Threonine	2.540 g	1.588 g	3.175 g	4.763 g
Tryptophan	0.800 g	0.500 g	1.000 g	1.500 g
Valine	3.604 g	2.253 g	4.505 g	6.758 g
Arginine	3.780 g	2.363 g	4.725 g	7.088 g
Histidine	1.752 g	1.095 g	2.190 g	3.285 g
Alanine	6.792 g	4.245 g	8.490 g	12.73 g
Aspartic acid	2.100 g	1.313 g	2.625 g	3.938 g
Glutamic acid	4.908 g	3.068 g	6.135 g	9.203 g
Glycine	2.312 g	1.445 g	2.890 g	4.335 g
Proline	4.760 g	2.975 g	5.950 g	8.925 g
Serine	4.200 g	2.625 g	5.250 g	7.875 g

	in 1000 ml	in 625 ml	in 1250 ml	in 1875 ml
Amino acid content [g]	56.0	35.0	70.1	105.1
Nitrogen content [g]	8	5	10	15
Carbohydrate content [g]	144	90	180	270
Lipid content [g]	40	25	50	75

	in 1000 ml	in 625 ml	in 1250 ml	in 1875 ml
Energy in the form of lipids [kJ (kcal)]	1590 (380)	995 (240)	1990 (475)	2985 (715)
Energy in the form of carbohydrates [kJ (kcal)]	2415 (575)	1510 (360)	3015 (720)	4520 (1080)
Energy in the form of amino acids [kJ (kcal)]	940 (225)	585 (140)	1170 (280)	1755 (420)
Non-protein energy [kJ (kcal)]	4005 (955)	2505 (600)	5005 (1195)	7510 (1795)
Total energy [kJ (kcal)]	4945 (1180)	3090 (740)	6175 (1475)	9265 (2215)

Osmolality [mOsm/kg]	1840
Theoretical osmolarity [mOsm/l]	1330
pH	5.0 - 6.0

The other ingredients are citric acid monohydrate (for pH adjustment), egg phospholipids for injection, glycerol, sodium oleate, all-rac-alpha-tocopherol, sodium hydroxide (for pH adjustment) and water for injections.

What Omeflex special without electrolytes looks like and contents of the pack

The ready-to-use product is an emulsion for infusion, i.e. it is administered through a small tube into a vein.

Omeflex special without electrolytes is supplied in flexible multichamber bags containing:

- 625 ml (250 ml of amino acids solution + 125 ml of fat emulsion + 250 ml of glucose solution)
- 1250 ml (500 ml of amino acids solution + 250 ml of fat emulsion + 500 ml of glucose solution)
- 1875 ml (750 ml of amino acids solution + 375 ml of fat emulsion + 750 ml of glucose solution)




Figure A: The multichamber bag is packed in a protective overwrap. An oxygen absorber and an oxygen indicator are placed between the bag and the overwrap; the oxygen absorber sachet is made of inert material and contains iron hydroxide.

Figure B: The top chamber contains a glucose solution, the middle chamber contains a fat emulsion, and the bottom chamber contains an amino acid solution.

The glucose and the amino acid solutions are clear and colourless up to straw-coloured. The fat emulsion is milky white.

The top chamber and the middle chamber can be connected with the bottom chamber by opening the intermediate seams.

The different container sizes are presented in cartons containing five bags.

Pack sizes: 5 x 625 ml, 5 x 1250 ml, and 5 x 1875 ml

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

B. Braun Melsungen AG

Carl-Braun-Straße 1

34212 Melsungen, Germany

Postal address:

34209 Melsungen, Germany

Phone: +49-5661-71-0

Fax: +49-5661-71-4567

This medicine is authorised in the Member States of the European Economic Area and in the United Kingdom (GB and NI) under the following names:

Austria

NuTRiflex Omega special ohne Elektrolyte B.Braun Emulsion zur Infusion

Belgium	Nutriflex Omega special zonder elektrolyten, 56 g/l Amino + 144 g/l G, emulsie voor infusie,
Czech Republic	Nutriflex Omega special bez elektrolýtů
Denmark	Nutriflex Omega Special elektrolytfri
France	REANUTRIFLEX OMEGA, émulsion pour perfusion
Germany	NuTRIflex Omega special ohne Elektrolyte novo Emulsion zur Infusion
Greece	Nutriflex Omega special without electrolytes
Ireland	Omeflex special without electrolytes emulsion for infusion
Italy	Omegefex AA38/G120 senza elettroliti
Luxembourg	NuTRIflex Omega special ohne Elektrolyte novo Emulsion zur Infusion
Netherlands	Nutriflex Omega special zonder elektrolyten, 56 g/l Amino + 144 g/l G, emulsie voor infusie
Norway	Nutriflex Omega Special elektrolytfri infusjonsvæske, emulsjon
Poland	Omegefex special bez elektrolitów
Portugal	Nutriflex Omega S sem eletrólitos emulsão para perfusão
Slovakia	Nutriflex Omega special bez elektrolýtov
Spain	Omegefex especial sin electrolitos Emulsión para perfusión
Sweden	Nutriflex Omega 56/144/40 elektrolytfri infusionsvätska, emulsion
United Kingdom (GB and NI)	Omeflex special without electrolytes emulsion for infusion

This leaflet was last revised in 07/2024

The following information is intended for healthcare professionals only:

Parenteral nutrition products should be visually inspected for damage, discolouration and emulsion instability before use.

Do not use bags which are damaged. Overwrap, primary bag and the peel seams between the chambers should be intact. Only use if the amino acid and glucose solutions are clear and colourless up to straw-coloured and the lipid emulsion is homogenous with milky white appearance. Do not use if the solutions contain particulate matter.


After mixing the three chambers, do not use if the emulsion shows discolouration or signs of phase separation (oil drops, oil layer). Stop the infusion immediately in case of discolouration of the emulsion or signs of phase separation.

Before opening the overwrap, check the colour of the oxygen indicator (see Figure A). Do not use if the oxygen indicator turned pink. Use only if the oxygen indicator is yellow.

Preparation of the mixed emulsion

Strict adherence to aseptic handling principles must be complied with.


To open: Tear overwrap starting from the tear notches (Fig. 1). Remove the bag from its protective overwrap. Discard overwrap, oxygen indicator and oxygen absorber.




Visually inspect the primary bag for leaks. Leaky bags must be discarded, since the sterility cannot be guaranteed.

Mixing of the bag and addition of additives


To open and mix the chambers sequentially, roll the bag with both hands, starting first by opening the peel seam that separates the top chamber (glucose) and the bottom chamber (amino acids) (Fig. 2).



Then continue applying pressure so that the peel seam separating the middle chamber (lipids) and the bottom chamber opens (Fig. 3).



After all chambers are mixed and following the removal of the aluminium seal (Fig. 3A), one can add compatible additives via the medication port (Fig. 4). Mix the contents thoroughly (Fig. 5) and visually inspect the mixture (Fig. 6). The mixture is a milky white homogenous oil-in-water emulsion. There should be no signs of emulsion phase separation.



Omeflex special without electrolytes can be mixed with the following additives up to the below specified upper concentration limits or maximum amount of additives after supplementation. The resulting admixtures are stable for 7 days between +2 °C to +8 °C plus 2 days at 25 °C.

- Electrolytes: stability has been demonstrated up to a total quantity of 200 mmol/l of sodium + potassium (sum), 9.6 mmol/l of magnesium and 6.4 mmol/l of calcium in the ternary mixture.
- Phosphate: stability has been demonstrated up to a maximum concentration of 20 mmol/l for inorganic phosphate or up to a maximum concentration of 30 mmol/l for organic phosphate (not both at the same time).
- Alanyl-Glutamin up to 24 g/l.
- Trace elements and vitamins: stability has been demonstrated with commercially available multi-trace elements and multi-vitamins (e.g. Tracutil, Cernevit) up to the standard dosage recommended by the respective manufacturer of the micronutrient.

Detailed information about the above mentioned additives and the corresponding shelf life of such admixtures can be provided on demand by the manufacturer.

Preparation for infusion

The emulsion should always be brought to room temperature prior to infusion.

Remove the aluminium foil from the infusion port (Fig. 7) and attach the infusion set (Fig. 8). Use a non-vented infusion set or close the air vent when using a vented set. Hang the bag on an infusion stand (Fig. 9) and carry out infusion using the standard technique.



For single use only. Container and unused residues must be discarded after use.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Do not reconnect partially used containers.

If filters are used they must be lipid-permeable (pore size $\geq 1.2 \mu\text{m}$).

Shelf life after removing the protective overwrap and after mixing of the contents of the bag

Chemical and physicochemical in-use stability of the mixture of amino acids, glucose and fat was demonstrated for 7 days at 2-8 °C and additional 2 days at 25 °C.

Shelf life after admixture of compatible additives

From a microbiological point of view, the product should be used immediately after admixture of additives. If not used immediately after admixture of additives, in-use storage times and conditions prior to use are the responsibility of the user.

After first opening (spiking of the infusion port)

The emulsion is to be used immediately after opening of the container.

Omeflex special without electrolytes must not be mixed with other medicinal products for which compatibility has not been documented.

Omeflex special without electrolytes should not be given simultaneously with blood in the same infusion set due to the risk of pseudoagglutination.